



Ammar Anbari, MD
Shoulder, Hip and Knee Arthroscopy
and Reconstruction
Office Phone: (860) 889-7345

Multi-directional Instability Repair Prescription

Name _____

Date _____

Diagnosis _____

Date of Surgery _____

Frequency: 1 2 3 4 times/week Duration: 1 2 3 4 5 6 Weeks

_____ Weeks 0-6:

- Gunslinger Brace
- Isometrics in brace
- Grip strengthening

_____ Weeks 6-12:

- Sling at night
- AROM only as tolerated to increase ROM; no PT stretching or manipulation
- Restrict to 140° FF/ 40° ER at side/ IR to stomach/ 45° Abduction
- Begin light isometrics for rotator cuff and deltoid, with arm at the side

_____ Months 3-12:

- Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 set per exercise for rotator cuff, deltoid, and scapular stabilizers
- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- If ROM lacking, increase to full with gentle passive stretching at end ranges
- Begin eccentric motions, plyometrics (ex. Weighted ball toss), and closed chain exercises at 16 weeks
- Begin sports related rehab at 4 ½ months, including advanced conditioning
- Return to throwing at 6 months
- Throw from pitcher's mound at 9 months
- No collision sports allowed
- MMI is usually at 12 months

Modalities

Electric Stimulation

Ultrasound

Signature: _____

Ammar Anbari, MD